# Tgfb1 Rabbit pAb

Catalog No.: A23262



## **Basic Information**

### **Observed MW**

44kDa

#### **Calculated MW**

44kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Mouse, Rat

# **Background**

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGF-beta family members. This encoded protein regulates cell proliferation, differentiation and growth, and can modulate expression and activation of other growth factors including interferon gamma and tumor necrosis factor alpha. Mice lacking a functional copy of this gene develop severe multifocal inflammatory disease, yolk sac defects and colon cancer.

# **Recommended Dilutions**

**WB** 

1:500 - 1:1000

# **Immunogen Information**

**Gene ID** 21803

**Swiss Prot** 

P04202

Immunogen
Recombinant protein of mouse Tgfb1.

**Synonyms** 

Tgfb; Tgfb-1; TGFbeta1; TGF-beta1; Tgfb1

# **Contact**

| <b>a</b>  |   | 400-999-6126              |
|-----------|---|---------------------------|
| $\bowtie$ |   | cn.market@abclonal.com.cn |
| $\odot$   | Ī | www.abclonal.com.cn       |


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates, using Tgfb1 Rabbit pAb (A23262) at 1:2500 dilution. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.