ABclonal www.abclonal.com

ABflo® 594 Rabbit anti-Human CAR/CXADR mAb

Catalog No.: A23368

Basic Information

Observed MW

Calculated MW

40kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC58165-ABf594

Conjugate

ABflo® 594. Ex:588nm. Em:604nm.

Background

The protein encoded by this gene is a type I membrane receptor for group B coxsackieviruses and subgroup C adenoviruses. Several transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene are found on chromosomes 15, 18, and 21.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 1525 **Swiss Prot**

P78310

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 20-237 of human CAR/CXADR(NP_001329.1)

Synonyms

CAR; HCAR; CAR4/6

Contact

| <u>a</u> | | 400-999-6126 |
|-------------------|---|---------------------------|
| \bowtie | | cn.market@abclonal.com.cn |
| $\overline{\Box}$ | Т | www abclonal com cn |

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data





Flow cytometry:1X10^6 K-562 cells(negative control,left) and HepG2 cells (right) were surface-stained with ABflo® 594 Rabbit anti-Human CAR/CXADR mAb(A23368,5 µl/Test,orange line) or ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,blue line).Non-fluorescently stained cells was used as blank control (red line).

Flow cytometry:1X10^6 HepG2 cells were surface-stained with ABflo® 594 Rabbit IgG isotype control (5 µl/Test,left) or ABflo® 594 Rabbit anti-Human CAR/CXADR mAb(A23368,5 µl/Test,right).