

Catalog No.: A2343 5 Publications



Basic Information

Observed MW 95kDa

Calculated MW 90kDa

Category Primary antibody

Applications WB,ELISA

Cross-Reactivity Human, Mouse

Background

This gene encodes a member of a subfamily of RING-finger type E3 ubiquitin ligases. The protein binds to specific DNA sequences, and recruits a histone deacetylase to regulate gene expression. Its expression peaks at late G1 phase and continues during G2 and M phases of the cell cycle. It plays a major role in the G1/S transition by regulating topoisomerase llalpha and retinoblastoma gene expression, and functions in the p53-dependent DNA damage checkpoint. It is regarded as a hub protein for the integration of epigenetic information. This gene is up-regulated in various cancers, and it is therefore considered to be a therapeutic target. Multiple transcript variants encoding different isoforms have been found for this gene. A related pseudogene exists on chromosome 12.

Recommended Dilutions

WB	1:100 - 1:500
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Immunogen Information

Gene ID 29128

Swiss Prot Q96T88

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human UHRF1 (NP_001041666.1).

Synonyms

Np95; hNP95; ICBP90; RNF106; TDRD22; hUHRF1; huNp95; UHRF1

Contact


6	400-999-6126
\times	cn.market@abclonal.com.cn
Ð	www.abclonal.com.cn

Product Information

Source Rabbit **Isotype** IgG **Purification** Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of lysates from Mouse brain, using UHRF1 Rabbit pAb (A2343) at 1:500 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 90s.