# **Uhrf2 Rabbit pAb**

Catalog No.: A2344 1 Publications



## **Basic Information**

#### **Observed MW**

95kDa

### **Calculated MW**

90kDa

### Category

Primary antibody

### **Applications**

ELISA,WB

### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

Enables histone binding activity. Predicted to be involved in several processes, including maintenance of DNA methylation; protein autoubiquitination; and regulation of cell cycle. Located in heterochromatin and nucleus. Is expressed in several structures, including central nervous system; ear; early conceptus; female reproductive system; and urinary system. Orthologous to human UHRF2 (ubiquitin like with PHD and ring finger domains 2).

## **Recommended Dilutions**

**WB** 

1:500 - 1:1000

## **Immunogen Information**

**Gene ID** 109113

Swiss Prot

Q7TMI3

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 76-359 of mouse Uhrf2 (NP\_659122.2).

### **Synonyms**

Nirf; 2310065A22Rik; D130071B19Rik; Uhrf2

### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	T	www.abclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of Mouse thymus, using Uhrf2 antibody (A2344) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.