# **Bcl-XL Rabbit mAb**

Catalog No.: A23481 Recombinant



# **Basic Information**

#### **Observed MW**

30 kDa

### **Calculated MW**

26 kDa

### Category

Primary antibody

### **Applications**

WB,IHC-P,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

#### CloneNo number

ARC52304

# **Background**

The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. The proteins encoded by this gene are located at the outer mitochondrial membrane, and have been shown to regulate outer mitochondrial membrane channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Alternative splicing results in multiple transcript variants encoding two different isoforms. The longer isoform acts as an apoptotic inhibitor and the shorter isoform acts as an apoptotic activator.

### **Recommended Dilutions**

**WB** 1:1000 - 1:5000

IHC-P 1:50-1:200

**ELISA** Recommended starting concentration is 1 μg/mL.

Please optimize the concentration based on your specific assay requirements.

# Immunogen Information

**Gene ID**598
Swiss Prot
Q07817

#### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

BCLX; BCL2L; Bcl-X; PPP1R52; BCL-XL/S; Bcl-XL

### **Contact**

6		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
<u> </u>	ī	www.ahclonal.com.cn


### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of lysates from Hep G2 cells using Bcl-XL Rabbit mAb (A23481) at 1:2000 dilution incubated at room temperature for 1.5 hours.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25  $\mu g$  per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45 s.



Western blot analysis of lysates from Rat thymus using Bcl-XL Rabbit mAb (A23481) at 1:2000 dilution incubated overnight at  $4^{\circ}$ C.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 μg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30 s.



Western blot analysis of lysates from Mouse liver using Bcl-XL Rabbit mAb (A23481) at 1:2000 dilution incubated overnight at  $4^{\circ}$ C.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 45 s.

# **Validation Data**



Immunohistochemistry analysis of paraffinembedded Rat testis tissue using BcI-XL Rabbit mAb (A23481) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.