

# MMP9 Rabbit mAb

Catalog No.: A23535

Recombinant

4 Publications

## Basic Information

**Observed MW**

100kDa

**Calculated MW**

78kDa

**Category**

Primary antibody

**Applications**

WB,FC (intra),ELISA

**Cross-Reactivity**

Human

**CloneNo number**

ARC60372

## Background

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades type IV and V collagens. Studies in rhesus monkeys suggest that the enzyme is involved in IL-8-induced mobilization of hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-associated tissue remodeling.

## Recommended Dilutions

**WB** 1:1000 - 1:4000**FC (intra)** 1:500 - 1:1000

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Immunogen Information

**Gene ID**

4318

**Swiss Prot**

P14780

**Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

**Synonyms**

GELB; CLG4B; MMP-9; MANDP2; MMP9

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of lysates from wild type (WT) and 293T cells transfected with MMP9 using MMP9 Rabbit mAb (A23535) at 1:1000 dilution.



Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 30s.



Flow cytometry:  $1 \times 10^6$  293F cells (Low Expression, left) and 293F (Transfection, right) cells were intracellularly-stained with MMP9 Rabbit mAb (A23535, 2 µg/mL, orange line) or ABflo® 594 Rabbit IgG isotype control (A23821, 5 µl/Test, blue line), followed by ABflo® 594-conjugated Goat Anti-Rabbit IgG (H+L) staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1 \times 10^6$  293F (Transfection, right) cells were intracellularly-stained with ABflo® 594 Rabbit IgG isotype control (A23821, 5 µl/Test, left) or MMP9 Rabbit mAb (A23535, 2 µg/mL, right).