

PCSK9 Rabbit pAb

Catalog No.: A23546

Basic Information

Observed MW

Calculated MW

74KDa

Category

Primary antibody

Applications

ELISA,WB,IF/ICC

Cross-Reactivity

Human, Mouse, Rat

Background

Enables apolipoprotein binding activity; lipoprotein particle binding activity; and low-density lipoprotein particle receptor binding activity. Involved in several processes, including cellular response to insulin stimulus; cellular response to starvation; and regulation of neuron apoptotic process. Acts upstream of or within several processes, including low-density lipoprotein receptor particle metabolic process; regulation of low-density lipoprotein particle receptor catabolic process; and triglyceride metabolic process. Located in COPII-coated ER to Golgi transport vesicle; endoplasmic reticulum; and extracellular space. Is expressed in several structures, including alimentary system; cerebellum; genitourinary system; liver; and telencephalon. Human ortholog(s) of this gene implicated in familial hypercholesterolemia and hypobetalipoproteinemia. Orthologous to human PCSK9 (proprotein convertase subtilisin/kexin type 9).

Recommended Dilutions

WB 1:500 - 1:1000

IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID

100102

Swiss Prot

Q80W65

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 35-694 of mouse PCSK9. (NP_705793.1).

Synonyms

FH3; PC9; Narc1; HCHOLA3; PCSK9

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.


Validation Data



Western blot analysis of lysates from Mouse liver, using PCSK9 Rabbit pAb (A23546) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 60s.



Immunofluorescence analysis of HeLa cells using PCSK9 Rabbit pAb(A23546) at a dilution of 1:200 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using PCSK9 Rabbit pAb(A23546) at a dilution of 1:200 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L)(AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.