

PE Rabbit anti-Human DNAM-1/CD226 mAb

Catalog No.: A23591

Basic Information

Observed MW

Calculated MW

39kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC60058-PE

Conjugate

PE. Ex:565nm. Em:574nm.

Background

This gene encodes a glycoprotein expressed on the surface of NK cells, platelets, monocytes and a subset of T cells. It is a member of the Ig-superfamily containing 2 Ig-like domains of the V-set. The protein mediates cellular adhesion of platelets and megakaryocytic cells to vascular endothelial cells. The protein also plays a role in megakaryocytic cell maturation. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Immunogen Information

Gene ID

10666

Swiss Prot

Q15762

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 19-247 of human DNAM-1/CD226 (NP_006557.2).

Synonyms

PTA1; DNAM1; DNAM-1; TLISA1

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data



Flow cytometry: 1×10^6 293F cells (negative control, left) and HEL cells (right) were surface-stained with PE Rabbit anti-Human DNAM-1/CD226 mAb (A23591, 5 μ l/Test, orange line) or PE Rabbit IgG isotype control (5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 HEL cells were surface-stained with PE Rabbit IgG isotype control (5 μ l/Test, left) or PE Rabbit anti-Human DNAM-1/CD226 mAb (A23591, 5 μ l/Test, right).