

TriMethyl-Histone H3-K9 Rabbit pAb

Catalog No.: A2360

60 Publications

Basic Information

Observed MW

17 kDa

Calculated MW

15 kDa

Category

Primary antibody

Applications

WB,DB,IHC-P,IF/ICC,ELISA,CHIP

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

Recommended Dilutions

WB	1:100 - 1:500
-----------	---------------

DB	1:500 - 1:2000
-----------	----------------

IHC-P	1:50 - 1:200
--------------	--------------

IF/ICC	1:50 - 1:200
---------------	--------------

ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
--------------	---

CHIP	2.5µg antibody for 5µg-10µg of Chromatin
-------------	--

Contact

	400-999-6126
--	--------------

	cn.market@abclonal.com.cn
--	---------------------------

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; TriMethyl-Histone H3-K9

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH7.3.




| www.abclonal.com.cn


Validation Data




Western blot analysis of various lysates using TriMethyl-Histone H3-K9 Rabbit pAb (A2360) at 1:500 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 90s.




Chromatin immunoprecipitation analysis of extracts of HeLa cells, using TriMethyl-Histone H3-K9 antibody (A2360) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.



Dot-blot analysis of all sorts of methylation peptides using TriMethyl-Histone H3-K9 antibody (A2360).




Immunofluorescence analysis of HeLa cells using TriMethyl-Histone H3-K9 Rabbit pAb (A2360) at dilution of 1:50 (40x lens).
 Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of MCF7 cells using TriMethyl-Histone H3-K9 Rabbit pAb (A2360) at dilution of 1:50 (40x lens).
 Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.


Validation Data




Immunofluorescence analysis of NIH/3T3 cells using TriMethyl-Histone H3-K9 Rabbit pAb (A2360) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using TriMethyl-Histone H3-K9 Rabbit pAb (A2360) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis using TriMethyl-Histone H3-K9 Rabbit pAb (A2360) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis using TriMethyl-Histone H3-K9 Rabbit pAb (A2360) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.