

DiMethyl-Histone H3-K36 Rabbit pAb

Catalog No.: A2365

8 Publications

Basic Information

Observed MW

17 kDa

Calculated MW

15 kDa

Category

Primary antibody

Applications

WB, IHC-P, IF/ICC, ELISA, ChIP, ChIP-seq

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

Recommended Dilutions

WB	1:500 - 1:1000
-----------	----------------

IHC-P	1:50 - 1:200
--------------	--------------

IF/ICC	1:50 - 1:200
---------------	--------------

ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
--------------	---

ChIP	5µg antibody for 5µg-10µg of Chromatin
-------------	--

ChIP-seq	1:20 - 1:100
-----------------	--------------

Contact

		400-999-6126
--	--	--------------

		cn.market@abclonal.com.cn
--	--	--

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; DiMethyl-Histone H3-K36

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




| www.abclonal.com.cn


Validation Data



Western blot analysis of lysates from HeLa cells, using DiMethyl-Histone H3-K36 Rabbit pAb (A2365) at 1:600 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 180s.




Western blot analysis of various lysates, using DiMethyl-Histone H3-K36 Rabbit pAb (A2365) at 1:600 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Enhanced Kit (RM00021).
 Exposure time: 180s.




Chromatin immunoprecipitation analysis of γ -actin gene from 293 cell line, using DiMethyl-Histone H3-K36 antibody (A2365) and rabbit IgG. P1, P2 and P3 were probes located on γ -actin gene as the schematic diagram illustrated. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.


Validation Data




Dot-blot analysis of all sorts of methylation peptides using DiMethyl-Histone H3-K36 antibody (A2365).




Immunofluorescence analysis of 293T cells using DiMethyl-Histone H3-K36 Rabbit pAb (A2365). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Human breast using DiMethyl-Histone H3-K36 Rabbit pAb (A2365) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis using DiMethyl-Histone H3-K36 Rabbit pAb (A2365) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis using DiMethyl-Histone H3-K36 Rabbit pAb (A2365) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.