

# CD146 Rabbit pAb

Catalog No.: A23669

## Basic Information

### Observed MW

120kDa

### Calculated MW

71kDa

### Category

Primary antibody

### Applications

ELISA, WB

### Cross-Reactivity

Human, Mouse

## Background

Predicted to be involved in angiogenesis and positive regulation of cell migration. Predicted to act upstream of or within heterophilic cell-cell adhesion via plasma membrane cell adhesion molecules. Predicted to be located in external side of plasma membrane; extracellular space; and nucleus. Predicted to be active in plasma membrane. Is expressed in several structures, including adipose tissue; genitourinary system; gut; nervous system; and respiratory system. Orthologous to human MCAM (melanoma cell adhesion molecule).

## Recommended Dilutions

WB 1:500 - 1:1000

## Immunogen Information

### Gene ID

84004

### Swiss Prot

Q8R2Y2

### Immunogen

Recombinant protein of mouse CD146.

### Synonyms

CD146; CD149; Muc18; s-endo; 1-gicerin; s-gicerin

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of various lysates, using CD146 Rabbit pAb (A23669) at 1:800 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 45s.