MonoMethyl-Histone H3-K79 Rabbit pAb

ABclonal

www.abclonal.com

Catalog No.: A2367 3 Publications

Basic Information

Observed MW

17 kDa

Calculated MW

15 kDa

Category

Primary antibody

Applications

WB,IHC-P,IF/ICC,ELISA,ChIP,ChIP-seq

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB 1:500 - 1:1000

1:50 - 1:200 **IHC-P**

1:50 - 1:200 IF/ICC

Recommended starting **ELISA**

> concentration is 1 µg/mL. Please optimize the concentration based on your specific assay

requirements.

ChIP 6µg antibody for

5μg-10μg of Chromatin

1:20 - 1:100 ChIP-seq

Contact

<u>a</u>		400-999-6126
\bowtie	Π	cn.market@abclonal.com.cn

Immunogen Information

Gene ID Swiss Prot 8290/8350 Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; MonoMethyl-Histone H3-K79


Product Information

Source Isotype **Purification** Rabbit IgG Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of various lysates using MonoMethyl-Histone H3-K79 Rabbit pAb (A2367) at 1:1000 dilution


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).

Exposure time: 300s.



Chromatin immunoprecipitation analysis of extracts of HeLa cells, using MonoMethyl-Histone H3-K79 antibody (A2367) and rabbit IgG.The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.





Dot-blot analysis of all sorts of methylation peptides using MonoMethyl-Histone H3-K79 antibody (A2367).

Immunofluorescence analysis of 293T cells using MonoMethyl-Histone H3-K79 Rabbit pAb (A2367). Blue: DAPI for nuclear staining.