

# DiMethyl-Histone H3-K79 Rabbit pAb

Catalog No.: A2368

3 Publications

## Basic Information

### Observed MW

17kDa

### Calculated MW

16kDa

### Category

Primary antibody

### Applications

WB, IHC-P, IF/ICC, IP, ELISA, ChIP, ChIP-seq

### Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

## Recommended Dilutions

<b>WB</b>	1:500 - 1:1000
-----------	----------------

<b>IHC-P</b>	1:50 - 1:100
--------------	--------------

<b>IF/ICC</b>	1:50 - 1:200
---------------	--------------

<b>IP</b>	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells
-----------	--

<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
--------------	--

<b>ChIP</b>	5µg antibody for 5µg-10µg of Chromatin
-------------	---

<b>ChIP-seq</b>	1:50 - 1:200
-----------------	--------------

## Immunogen Information

### Gene ID

8290/8350

### Swiss Prot

Q16695/P68431

### Immunogen

A synthetic dimethylated peptide around K79 of human Histone H3 (NP\_003520.1).

### Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; DiMethyl-Histone H3-K79

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification

Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

## Contact

---


 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)


 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

---

## Validation Data



Immunoprecipitation analysis of 300 µg extracts of HeLa cells using 3 µg DiMethyl-Histone H3-K79 antibody (A2368). Western blot was performed from the immunoprecipitate using DiMethyl-Histone H3-K79 antibody (A2368) at a dilution of 1:1000.




Western blot analysis of lysates from HeLa cells, using DiMethyl-Histone H3-K79 Rabbit pAb (A2368) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.




Chromatin immunoprecipitations were performed with cross-linked chromatin from K-562 cells and DiMethyl-Histone H3-K79 Rabbit pAb (A2368). The ChIP sequencing results indicate the enrichment pattern of DiMethyl-Histone H3-K79 in selected genomic region and representative gene loci (GAPDH), as shown in figure.




Chromatin immunoprecipitation analysis of extracts of MCF7 cells, using DiMethyl-Histone H3-K79 antibody (A2368) and rabbit IgG. The amount of immunoprecipitated DNA was checked by quantitative PCR. Histogram was constructed by the ratios of the immunoprecipitated DNA to the input.


## Validation Data




Dot-blot analysis of all sorts of methylation peptides using DiMethyl-Histone H3-K79 antibody (A2368).




Immunofluorescence analysis of 293T cells using DiMethyl-Histone H3-K79 Rabbit pAb (A2368). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded Human mammary cancer using DiMethyl-Histone H3-K79 Rabbit pAb (A2368) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis using DiMethyl-Histone H3-K79 Rabbit pAb (A2368) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse testis using DiMethyl-Histone H3-K79 Rabbit pAb (A2368) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.