Leader in Biomolecular Solutions for Life Science

# ABflo® 488 Rabbit anti-Mouse CD8a mAb

Catalog No.: A23903



# **Basic Information**

**Observed MW** Refer to figures

Calculated MW 27KDa

Category Primary antibody

Applications FC

Cross-Reactivity Mouse

CloneNo number ARC60677

#### Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

#### **Recommended Dilutions**

5 µl per 10^6 cells in

100 µl volume

FC

# Immunogen Information

Gene ID 12525 Swiss Prot P01731

#### Immunogen

Background

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Enables identical protein binding activity. Acts upstream of or within several processes, including cytotoxic T cell differentiation; defense response to virus; and positive regulation of

and hemolymphoid system gland. Orthologous to human CD8A (CD8a molecule).

calcium-mediated signaling. Located in external side of plasma membrane. Is expressed in several structures, including endocrine gland; exocrine gland; genitourinary system; gut;

#### Synonyms

Ly-2; Ly-B; Ly-35; Lyt-2

### Contact

6	400-999-6126
$\times$	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

# **Product Information**

**Source** Rabbit **lsotype** IgG **Purification** Affinity purification

#### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

# Validation Data





Flow cytometry: 1X10^6 Raw264.7 cells (negative control,left) and C57BL/6 mouse Splenocytes (right) were surface-stained with ABflo® 488 Rabbit anti-Mouse CD8a mAb (A23903,5 µl/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line). Flow cytometry:1X10^6 C57BL/6 mouse Splenocytes were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,left) or ABflo® 488 Rabbit anti-Mouse CD8a mAb(A23903,5 µl/Test,right).