

# IL9 Rabbit mAb

Catalog No.: A24007 **Recombinant**

## Basic Information

### Observed MW

19kDa

### Calculated MW

16kDa

### Category

Primary antibody

### Applications

ELISA,WB,FC (intra)

### Cross-Reactivity

Human

### CloneNo number

ARC62039

## Background

The protein encoded by this gene is a cytokine that acts as a regulator of a variety of hematopoietic cells. This cytokine stimulates cell proliferation and prevents apoptosis. It functions through the interleukin 9 receptor (IL9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. The gene encoding this cytokine has been identified as a candidate gene for asthma. Genetic studies on a mouse model of asthma demonstrated that this cytokine is a determining factor in the pathogenesis of bronchial hyperresponsiveness.

## Recommended Dilutions

<b>WB</b>	1:500 - 1:1000
<b>FC (intra)</b>	1:500 - 1:1000

## Immunogen Information

### Gene ID

3578

### Swiss Prot

P15248

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 19-144 of human IL9(NP\_000581.1).

### Synonyms

P40; HP40; IL-9; IL9

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage



Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of lysates from 293T(Transfection) and control 293T cells, using IL9 Rabbit mAb (A24007) at 1:1000 dilution.  
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25ug per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 30s.



$1 \times 10^6$  293T cells (negative control, left) and 293T(Transfection, right) cells were intracellularly-stained with IL9 Rabbit mAb(A24007, 2 $\mu$ g/mL, orange line) or ABflo® 488 Rabbit IgG isotype control (A22069, 2 $\mu$ g/mL, blue line), followed by FITC conjugated goat anti-rabbit pAb (1:200 dilution) staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1 \times 10^6$  293T(Transfection) cells were intracellularly-stained with ABflo® 488 Rabbit IgG isotype control (A22069, 2 $\mu$ g/mL, left) or IL9 Rabbit mAb(A24007, 2 $\mu$ g/mL, right).