# ABclonal www.abclonal.com

# ABflo® 594 Rabbit anti-Human CD48 mAb

Catalog No.: A24195

## **Basic Information**

#### **Observed MW**

# **Calculated MW**

19kDa/28kDa

## Category

Primary antibody

# **Applications**

FC

## **Cross-Reactivity**

Human

#### CloneNo number

ARC53039-ABflo594

# Conjugate

ABflo® 594. Ex:588nm. Em:604nm.

# **Background**

This gene encodes a member of the CD2 subfamily of immunoglobulin-like receptors which includes SLAM (signaling lymphocyte activation molecules) proteins. The encoded protein is found on the surface of lymphocytes and other immune cells, dendritic cells and endothelial cells, and participates in activation and differentiation pathways in these cells. The encoded protein does not have a transmembrane domain, however, but is held at the cell surface by a GPI anchor via a C-terminal domain which maybe cleaved to yield a soluble form of the receptor. Multiple transcript variants encoding different isoforms have been found for this gene.

# **Recommended Dilutions**

FC

5  $\mu$ l per 10^6 cells in 100  $\mu$ l volume

# **Immunogen Information**

Gene ID 962 **Swiss Prot** 

P09326

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 27-220 of human CD48 (NP\_001769.2)

## **Synonyms**

BCM1; BLAST; hCD48; mCD48; BLAST1; SLAMF2; MEM-102

## **Contact**

	400-999-6126
<b>×</b>	cn.market@abclonal.com.cn
	www.abclonal.com.cn

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification


## **Storage**

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

## **Validation Data**





Flow cytometry: 1X10^6 K-562 cells (negative control,left) and Daudi cells (right) were surface-stained with ABflo® 594 Rabbit anti-Human CD48 mAb (A24195,5 µl/Test,orange line) or ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 Daudi cells were surface-stained with ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,left) or ABflo® 594 Rabbit anti-Human CD48 mAb (A24195,5 µl/Test,right).