ABclonal www.abclonal.com

ABflo® 594 Rabbit anti-Human IL-1β mAb

Catalog No.: A24245

Basic Information

Observed MW

Calculated MW

30kDa

Category

Primary antibody

Applications

FC (intra)

Cross-Reactivity

Human

CloneNo number

ARC53391

Conjugate

ABflo® 594. Ex:588nm. Em:604nm.

Background

The protein encoded by this gene is a member of the interleukin 1 cytokine family. This cytokine is produced by activated macrophages as a proprotein, which is proteolytically processed to its active form by caspase 1 (CASP1/ICE). This cytokine is an important mediator of the inflammatory response, and is involved in a variety of cellular activities, including cell proliferation, differentiation, and apoptosis. The induction of cyclooxygenase-2 (PTGS2/COX2) by this cytokine in the central nervous system (CNS) is found to contribute to inflammatory pain hypersensitivity. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2.

Recommended Dilutions

FC (intra)

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 3553 **Swiss Prot**

P01584

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

IL1B; IL-1; IL1-BETA; IL1F2; interleukin-1 beta

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
\odot	www.abclonal.com.cn

Product Information


Source	Isotype	Purification
Rabbit	IgG	Affinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data





Flow cytometry: $1X10^6$ 293T cells(negative control,left) and 293T (Transfection,right) were intracellularly-stained with ABflo® 594 Rabbit anti-Human IL-1 β mAb(A24245,5 μ I/Test,orange line) or ABflo® 594 Rabbit IgG isotype control (A23821,5 μ I/Test,blue line).Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:1X10^6 293T(Transfection) cells were intracellularly-stained with ABflo® 594 Rabbit IgG isotype control (A23821,5 μ I/Test,left) or ABflo® 594 Rabbit anti-Human IL-1 β mAb(A24245,5 μ I/Test,right).