ABflo® 594 Rabbit anti-Human CD47 mAb

ABclonal www.abclonal.com

Catalog No.: A24302

Basic Information

Observed MW

Refer to figures

Calculated MW

31kDa/33kDa/35kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC61212

Conjugate

ABflo® 594. Ex:588nm. Em:604nm.

Background

This gene encodes a membrane protein, which is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. The encoded protein is also a receptor for the C-terminal cell binding domain of thrombospondin, and it may play a role in membrane transport and signal transduction. This gene has broad tissue distribution, and is reduced in expression on Rh erythrocytes. Alternatively spliced transcript variants have been found for this gene.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 961 **Swiss Prot**

Q08722

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

CD47; IAP; MER6; OA3; CD47 molecule

Contact

a	400-999-6126
\bowtie	cn.market@abclonal.com.cn
\odot	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data









Flow cytometry:1X10^6 HepG2 cells (Low Expression, Left) and MOLT-4 cells(Right) were surface-stained with ABflo® 594 Rabbit anti-Human CD47 mAb(A24302,5 µl/Test,orange line) or ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:1X10^6 MOLT-4 cells were surface-stained with ABflo® 594 Rabbit IgG isotype control (A23821,5 μ I/Test,left) or ABflo® 594 Rabbit anti-Human CD47 mAb(A24302,5 μ I/Test,right).