PARP1 Rabbit pAb

Catalog No.: A2432 5 Publications



Basic Information

Observed MW

80kDa

Calculated MW

113kDa

Category

Primary antibody

Applications

WB,IP

Cross-Reactivity

Human, Mouse, Rat, Monkey

Background

This gene encodes a chromatin-associated enzyme, poly(ADP-ribosyl)transferase, which modifies various nuclear proteins by poly(ADP-ribosyl)ation. The modification is dependent on DNA and is involved in the regulation of various important cellular processes such as differentiation, proliferation, and tumor transformation and also in the regulation of the molecular events involved in the recovery of cell from DNA damage. In addition, this enzyme may be the site of mutation in Fanconi anemia, and may participate in the pathophysiology of type I diabetes.

Recommended Dilutions

WB 1:500 - 1:2000

IP 1:20 - 1:50

Immunogen Information

Gene IDSwiss Prot
142
P09874

Immunogen

A synthetic peptide of human PARP1

Synonyms

PARP; PARS; PPOL; ADPRT; ARTD1; ADPRT1; PARP-1; ADPRT 1; pADPRT-1; Poly-PARP; PARP1

Contact

| a | | 400-999-6126 |
|----------------|---|---------------------------|
| \bowtie | | cn.market@abclonal.com.cn |
| \overline{a} | ı | www.ahclonal.com.cn |

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from THP-1 cells, using PARP1 Rabbit pAb (A2432). Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (ASO14) at 1:10000 dilution. Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.