

Acetyl-Histone H3-K4 Rabbit mAb

Catalog No.: A24341 **Recombinant**

Basic Information

Observed MW

17kDa/

Calculated MW

15kDa

Category

Primary antibody

Applications

WB, IP, IF/ICC, IHC-P, DB, ChIP, ELISA

Cross-Reactivity

Human, Mouse, Rat, Other (Wide Range Predicted)

CloneNo number

ARC62519

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilutions

WB 1:1000 - 1:3000**IP** 0.5µg-4µg antibody for
400µg-600µg extracts of
whole cells**IF/ICC** 1:200 - 1:2000**IHC-P** 1:50 - 1:200**DB** 1:2000 - 1:5000**ChIP** 5µg antibody for
5µg-10µg of Chromatin**ELISA** Recommended starting
concentration is 1 µg/mL.
Please optimize the
concentration based on
your specific assay
requirements.

Immunogen Information

Gene ID

8290/8350

Swiss Prot

Q16695/P68431

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

H3t; H3.4; H3/g; H3FT; H3C16; HIST3H3; Acetyl-Histone H3-K4

Product Information

Source

Rabbit

Isotype

IgG

Purification

Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Immunoprecipitation of Acetyl-Histone H3-K4 from 600 μ g extracts of HeLa cells treated with TSA (5mM ,16h) was performed using 3 μ g of Acetyl-Histone H3-K4 Rabbit mAb (A24341). Rabbit Control IgG (AC005) was used to precipitate the Control IgG sample. IP samples were eluted with 1X Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at a dilution of 1:500.






Western blot analysis of lysates from HeLa cells, using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at 1:1000 dilution. HeLa cells were treated with TSA (1 μ M) at 37°C for 18 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 20s.




Western blot analysis of lysates from NIH/3T3 cells, using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at 1:1000 dilution. NIH/3T3 cells were treated with TSA (1 μ M) at 37°C for 18 hours. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 20s.


Validation Data




Dot-blot analysis of all sorts of peptides using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at 1:1000 dilution.




Confocal imaging of HeLa cells (treated with TSA) and HeLa cells (untreated) using Acetyl-Histone H3-K4 Rabbit mAb (A24341, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of paraffin-embedded Human colon carcinoma using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human liver using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human testis using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Mouse testis using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat colon using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat kidney using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Rat spleen using Acetyl-Histone H3-K4 Rabbit mAb (A24341) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.