

CD68 Rabbit PolymAb®

Catalog No.: A24386

Basic Information

Observed MW

70-130kDa

Calculated MW

70-130kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P, IF/ICC

Cross-Reactivity

Human, Mouse

CloneNo number

ARC51158_ARC60987

Background

Involved in several processes, including cellular response to lipopolysaccharide; cellular response to oxidised low-density lipoprotein particle stimulus; and negative regulation of dendritic cell antigen processing and presentation. Acts upstream of or within aging and cellular response to organic substance. Located in lysosome and plasma membrane. Is expressed in several structures, including central nervous system; embryo mesenchyme; heart blood vessel; liver; and spleen. Orthologous to human CD68 (CD68 molecule).

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

968/ 12514

Swiss Prot

P34810/P31996

Immunogen

Recombinant protein of human/mouse CD68.

Synonyms

Lamp4; gp110; Scard1

Contact

☎	400-999-6126
✉	cn.market@abclonal.com.cn
🌐	www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates, using CD68 Rabbit PolymAb® (A24386) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Negative control (NC): Jurkat
 Exposure time: 10s.




Immunofluorescence analysis of RAW264.7 using CD68 Rabbit PolymAb® (A24386) at dilution of 1:200 (40x lens). Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded human colon using CD68 Rabbit PolymAb® (A24386) at dilution of 1:10000 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human liver using CD68 Rabbit PolymAb® (A24386) at dilution of 1:10000 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human lung using CD68 Rabbit PolymAb® (A24386) at dilution of 1:10000 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human tonsil using CD68 Rabbit PolymAb® (A24386) at dilution of 1:10000 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.