www.abclonal.com

ABclonal

ABflo® 594 Rabbit anti-Mouse CD146 mAb

Catalog No.: A24399

Basic Information

Observed MW

Refer to figures

Calculated MW

71kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC62356-ABflo594

Conjugate

ABflo® 594. Ex:588nm. Em:604nm.

Background

Predicted to be involved in angiogenesis and positive regulation of cell migration. Predicted to act upstream of or within heterophilic cell-cell adhesion via plasma membrane cell adhesion molecules. Predicted to be located in external side of plasma membrane; extracellular space; and nucleus. Predicted to be active in plasma membrane. Is expressed in several structures, including adipose tissue; genitourinary system; gut; nervous system; and respiratory system. Orthologous to human MCAM (melanoma cell adhesion molecule).

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID

Swiss Prot

84004

Q8R2Y2

Immunogen

Recombinant Protein corresponding to a sequence within amino acids 24-563 of mouse CD146 (NP_075548.2)

Synonyms

CD146; CD149; Muc18; s-endo; 1-gicerin; s-gicerin

Contact

a	400-999-6126
\bowtie	cn.market@abclonal.com.cn
\odot	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data





Flow cytometry:1X10^6 CTLL-2 cells (negative control,Left) and bEnd.3 cells (Right) were surface-stained with ABflo® 594 Rabbit anti-Mouse CD146 mAb(A24399,5 µl/Test,orange line) or ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:1X10^6 bEnd.3 cells were surface-stained with ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,left) or ABflo® 594 Rabbit anti-Mouse CD146 mAb(A24399,5 µl/Test,right).