

PCSK9 Rabbit pAb

Catalog No.: A24478

Basic Information

Observed MW

65kDa/80kDa/65kDa,80kDa

Calculated MW

20kDa/74kDa

Category

Primary antibody

Applications

ELISA,WB,IF/ICC,IP

Cross-Reactivity

Human, Rat

Background

This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an autocatalytic processing event with its prosegment in the ER and is constitutively secreted as an inactive protease into the extracellular matrix and trans-Golgi network. It is expressed in liver, intestine and kidney tissues and escorts specific receptors for lysosomal degradation. It plays a role in cholesterol and fatty acid metabolism. Mutations in this gene have been associated with autosomal dominant familial hypercholesterolemia. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:100 - 1:500

IF/ICC 1:50 - 1:200

IP 0.5µg-4µg antibody for
200µg-400µg extracts of
whole cells

Immunogen Information

Gene ID

255738

Swiss Prot

Q8NBP7

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 31-692 of human PCSK9 (NP_777596.2).

Synonyms

PCSK9; FH3; HCHOLA3; LDLCQ1; NARC-1; NARC1; PC9; proprotein convertase subtilisin/kexin type 9

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.


Validation Data



Western blot analysis of lysates from HeLa cells using PCSK9 Rabbit pAb(A24478) at 1:400 dilution. Secondary antibody:HRP Goat Anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection:ECL Basic Kit (RM00020). Exposure time:20s.



Immunoprecipitation analysis of 300ug extracts of HeLa cells using 3ug PCSK9 Rabbit pAb antibody (A24478 1:100). Western blot was performed from the immunoprecipitate using PCSK9 Rabbit pAb antibody (A24478) at a dilution of 1:1000.



Immunofluorescence analysis of HepG2 cells using PCSK9 Rabbit pAb (A24478) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.