

RNF126 Rabbit pAb

Catalog No.: A24515

Basic Information

Observed MW

36kDa

Calculated MW

33kDa/35kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P

Cross-Reactivity

Human, Mouse, Rat

Background

The protein encoded by this gene contains a RING finger domain, a motif present in a variety of functionally distinct proteins and known to be involved in protein-protein and protein-DNA interactions.

Recommended Dilutions

WB 1:1000 - 1:5000

IHC-P 1:100 - 1:500

Immunogen Information

Gene ID

55658

Swiss Prot

Q9BV68

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 120-200 of human RNF126 (NP_919442.1).

Synonyms

RNF126; ring finger protein 126

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.


Validation Data



Western blot analysis of various lysates, using RNF126 Rabbit pAb (A24515) at 1:2000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded human liver using RNF126 Rabbit pAb (A24515) at dilution of 1:300 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse liver using RNF126 Rabbit pAb (A24515) at dilution of 1:300 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.