

# ABflo® 594 Rabbit anti-Human CD62L/L-Selectin mAb

Catalog No.: A24632

### **Basic Information**

### **Observed MW**

Refer to figures

### **Calculated MW**

42kDa/43kDa

### Category

Primary antibody

### **Applications**

FC

### **Cross-Reactivity**

Human

### CloneNo number

ARC60035

### Conjugate

ABflo® 594. Ex:588nm. Em:604nm.

# **Background**

This gene encodes a cell surface adhesion molecule that belongs to a family of adhesion/homing receptors. The encoded protein contains a C-type lectin-like domain, a calcium-binding epidermal growth factor-like domain, and two short complement-like repeats. The gene product is required for binding and subsequent rolling of leucocytes on endothelial cells, facilitating their migration into secondary lymphoid organs and inflammation sites. Single-nucleotide polymorphisms in this gene have been associated with various diseases including immunoglobulin A nephropathy. Alternatively spliced transcript variants have been found for this gene.

### **Recommended Dilutions**

FC

5 μl per 10^6 cells in 100 μl volume

## **Immunogen Information**

Gene ID 6402 **Swiss Prot** 

P14151

### **Immunogen**

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

### **Synonyms**

SELL; CD62L; LAM1; LECAM1; LEU8; LNHR; LSEL; LYAM1; PLNHR; TQ1; L-selectin

### **Contact**

<b>a</b>	400-999-6126
$\bowtie$	cn.market@abclonal.com.cn
$\odot$	www.abclonal.com.cn

### **Product Information**


Source	Isotype	Purification
Rabbit	IgG	Affinity purification


#### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## **Validation Data**





Flow cytometry: 1X10^6 T-47D cells (negative control,left) and MOLT-4 cells (right) were surface-stained with ABflo® 594 Rabbit anti-Human CD62L mAb (A24632,5 µl/Test,orange line) or ABflo® 594 Rabbit IgG isotype control (A23821,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1X10^6$  MOLT-4 cells were surface-stained with ABflo® 594 Rabbit IgG isotype control (A23821,5  $\mu$ I/Test,left) or ABflo® 594 Rabbit anti-Human CD62L mAb (A24632,5  $\mu$ I/Test,right).