

ST6GAL1/CD75 Rabbit mAb

Catalog No.: A24645 **Recombinant**

Basic Information

Observed MW

50-70kDa

Calculated MW

20kDa/47kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P, IF/ICC, FC

Cross-Reactivity

Human

CloneNo number

ARC63359

Background

This gene encodes a member of glycosyltransferase family 29. The encoded protein is a type II membrane protein that catalyzes the transfer of sialic acid from CMP-sialic acid to galactose-containing substrates. The protein, which is normally found in the Golgi but can be proteolytically processed to a soluble form, is involved in the generation of the cell-surface carbohydrate determinants and differentiation antigens HB-6, CD75, and CD76. This gene has been incorrectly referred to as CD75. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB	1:2000 - 1:8000
IHC-P	1:500 - 1:1000
IF/ICC	1:50 - 1:200
FC	1:500 - 1:1000

Immunogen Information

Gene ID

6480

Swiss Prot

P15907

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 44-406 of human ST6GAL1/CD75 (NP_001340845.1).

Synonyms

ST6N; CDw75; SIAT1; ST6GalI; ST6GAL1/CD75

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from Hep G2 cells, using ST6GAL1/CD75 Rabbit mAb (A24645) at 1:7000 dilution.




Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 90s.



Flow cytometry: 1×10^6 HeLa cells (negative control, left) and Daudi cells (right) were surface-stained with ST6GAL1/CD75 Rabbit mAb (A24645, 2 $\mu\text{g}/\text{mL}$, orange line) or ABRlo® 647 Rabbit IgG isotype control (A22070, 2 $\mu\text{g}/\text{mL}$, blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 Daudi cells were surface-stained with ABRlo® 647 Rabbit IgG isotype control (A22070, 2 $\mu\text{g}/\text{mL}$, left) or ST6GAL1/CD75 Rabbit mAb (A24645, 2 $\mu\text{g}/\text{mL}$, right).

Confocal imaging of Daudi cells using ST6GAL1/CD75 Rabbit mAb (A24645, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunohistochemistry analysis of ST6GAL1/CD75 in paraffin-embedded Human liver using ST6GAL1/CD75 Rabbit mAb (A24645) at dilution of 1:800 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.