

Megalin/LRP2 Rabbit mAb

Catalog No.: A24690 Recombinant

Basic Information

Observed MW

250kDa+

Calculated MW

522kDa

Category

Primary antibody

Applications

ELISA,WB,IHC-P,IF/ICC,FC (intra)

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC63830

Background

The protein encoded by this gene, low density lipoprotein-related protein 2 (LRP2) or megalin, is a multi-ligand endocytic receptor that is expressed in many different tissues but primarily in absorptive epithelial tissues such as the kidney. This glycoprotein has a large amino-terminal extracellular domain, a single transmembrane domain, and a short carboxy-terminal cytoplasmic tail. The extracellular ligand-binding-domains bind diverse macromolecules including albumin, apolipoproteins B and E, and lipoprotein lipase. The LRP2 protein is critical for the reuptake of numerous ligands, including lipoproteins, sterols, vitamin-binding proteins, and hormones. This protein also has a role in cell-signaling; extracellular ligands include parathyroid hormones and the morphogen sonic hedgehog while cytosolic ligands include MAP kinase scaffold proteins and JNK interacting proteins. Recycling of this membrane receptor is regulated by phosphorylation of its cytoplasmic domain. Mutations in this gene cause Donnai-Barrow syndrome (DBS) and facio-oculoacoustico-renal syndrome (FOAR).

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200
FC (intra)	1:100 - 1:500

Immunogen Information

Gene ID

4036

Swiss Prot

P98164

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 4446-4655 of human Megalin/LRP2 [NP_004516.2].

Synonyms

DBS; GP330; LRP-2; Megalin/LRP2

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn
 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from Mouse kidney using Megalin/LRP2 Rabbit mAb(A24690) at 1:1000 dilution.




Secondary antibody:HRP Goat Anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection:ECL Basic Kit (RM00020).




Exposure time:20s.



Flow cytometry: 1×10^6 Reh cells (Low Expression, left) and Caco-2 (right) cells were intracellularly-stained with Megalin/LRP2 Rabbit mAb (A24690, 2 µg/mL, orange line) or ABflo® 647 Rabbit IgG isotype control (A22070, 5 µl/Test, blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 Caco-2 cells were intracellularly-stained with ABflo® 647 Rabbit IgG isotype control (A22070, 5 µl/Test, left) or Megalin/LRP2 Rabbit mAb (A24690, 2 µg/mL, right).

Confocal imaging of mouse kidney using Megalin/LRP2 Rabbit mAb (A24690, at dilution of 1:200) (Red). DAPI was used for nuclear staining (blue). Objective: 40x.




Confocal imaging of rat kidney using Megalin/LRP2 Rabbit mAb (A24690, at dilution of 1:200) (Red). DAPI was used for nuclear staining (blue). Objective: 40x.


Immunohistochemistry analysis of Megalin/LRP2 in paraffin-embedded human kidney using Megalin/LRP2 Rabbit mAb (A24690) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Immunohistochemistry analysis of Megalin/LRP2 in paraffin-embedded human liver using Megalin/LRP2 Rabbit mAb (A24690) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Validation Data



Immunohistochemistry analysis of Megalin/LRP2 in paraffin-embedded mouse kidney using Megalin/LRP2 Rabbit mAb (A24690) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Megalin/LRP2 in paraffin-embedded rat liver using Megalin/LRP2 Rabbit mAb (A24690) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.