

ABflo® 647 Rabbit anti-Human CD261/TRAIL-R1 mAb

Catalog No.: A24701

Basic Information

Observed MW

Calculated MW

50kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC63581

Conjugate

ABflo® 647. Ex:648nm. Em:664nm.

Background

The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice suggested that FADD, a death domain containing adaptor protein, is required for the apoptosis mediated by this protein.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 8797 **Swiss Prot**

000220

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

TNFRSF10A; APO2; CD261; DR4; TRAILR-1; TRAILR1; TNF receptor superfamily member 10a

Contact

2		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data





Flow cytometry: 1X10^6 BEWO cells (negative control,left) and HeLa cells (right) were surface-stained with ABflo® 647 Rabbit anti-Human CD261/TRAIL-R1 mAb (A24701,5 µl/Test,orange line) or ABflo® 647 Rabbit IgG isotype control (A22070,5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: $1X10^6$ HeLa cells were surface-stained with ABflo® 647 Rabbit IgG isotype control (A22070,5 μ I/Test,Ieft) or ABflo® 647 Rabbit anti-Human CD261/TRAIL-R1 mAb (A24701,5 μ I/Test,right).