

PE Rabbit anti-Mouse CD262/TRAILR2 mAb

Catalog No.: A24798

Basic Information

Observed MW

Calculated MW

42kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC62153-PE

Conjugate

PE. Ex:565nm. Em:574nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

Predicted to enable TRAIL receptor activity; identical protein binding activity; and protease binding activity. Predicted to be involved in TRAIL-activated apoptotic signaling pathway and positive regulation of apoptotic process. Predicted to act upstream of or within apoptotic process and regulation of apoptotic process. Predicted to be located in Golgi apparatus; cytosol; and membrane raft. Predicted to be active in cell surface and plasma membrane. Is expressed in bladder; liver; renal vasculature; urethra of female; and urethra of male. Human ortholog(s) of this gene implicated in carcinoma (multiple); cervical cancer; hematologic cancer (multiple); and urinary bladder cancer. Orthologous to several human genes including TNFRSF10A (TNF receptor superfamily member 10a).

Immunogen Information

Gene ID

21933

Swiss Prot

Q9QZM4

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 53-170 of mouse CD262/TRAILR2(NP_064671.2).

Synonyms

MK; DR5; Ly98; KILLER; TRICKB; TRAILR2; TRICK2A; TRICK2B

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification



Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data



Flow cytometry: 1×10^6 293F cells (negative control, left) and 293F (Transfection, right) cells were surface-stained with PE Rabbit anti-Mouse CD262/TRAILR2 mAb (A24798, 5 μ l/Test, orange line) or PE Rabbit IgG isotype control (5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 293F (Transfection) cells were surface-stained with PE Rabbit IgG isotype control (5 μ l/Test, left) or PE Rabbit anti-Mouse CD262/TRAILR2 mAb (A24798, 5 μ l/Test, right).