

IFIH1 Rabbit pAb

Catalog No.: A24868

Basic Information

Observed MW

140kDa

Calculated MW

25kDa/116kDa

Category

Primary antibody

Applications

ELISA,WB

Cross-Reactivity

Human

Background

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein that is upregulated in response to treatment with beta-interferon and a protein kinase C-activating compound, mezerein. Irreversible reprogramming of melanomas can be achieved by treatment with both these agents; treatment with either agent alone only achieves reversible differentiation. Genetic variation in this gene is associated with diabetes mellitus insulin-dependent type 19.

Recommended Dilutions

WB 1:500 - 1:1000

Immunogen Information

Gene ID

64135

Swiss Prot

Q9BYX4

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 277-490 of human IFIH1 (NP_071451.2).

Synonyms

IFIH1; AGS7; Hlcd; IDDM19; MDA-5; MDA5; RLR-2; SGMRT1; interferon induced with helicase C domain 1

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from THP-1 cells using IFIH1 Rabbit pAb (A24868) at 1:1000 dilution. THP-1 cells were treated by LPS (1 $\mu\text{g}/\text{ml}$) at 37°C for 24 hours.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25 μg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 5s.