ABflo® 488 Rabbit anti-Human CD98 mAb

Catalog No.: A24932



Basic Information

Observed MW

Calculated MW 57kDa/61kDa/67kDa/68kDa/71kDa

Category Primary antibody

Applications FC

Cross-Reactivity Human

CloneNo number ARC64398

Conjugate ABflo® 488. Ex:491nm. Em:516nm.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Background

This gene is a member of the solute carrier family and encodes a cell surface, transmembrane protein. The protein exists as the heavy chain of a heterodimer, covalently bound through di-sulfide bonds to one of several possible light chains. The encoded transporter plays a role in regulation of intracellular calcium levels and transports L-type amino acids. Alternatively spliced transcript variants, encoding different isoforms, have been characterized.

Immunogen Information

Gene ID 6520

Swiss Prot P08195

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

4F2; CD98; MDU1; 4F2HC; 4T2HC; NACAE; CD98HC

Contact

6	400-999-6126
\bowtie	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Product Information


Source Rabbit **lsotype** IgG **Purification** Affinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data





Flow cytometry: 1X10^6 SH-SY5Y cells (Low Expression,left) and HeLa cells (right) were surface-stained with ABflo® 488 Rabbit anti-Human CD98 mAb (A24932,5 μ /Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 μ /Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 HeLa cells were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,left) or ABflo® 488 Rabbit anti-Human CD98 mAb (A24932,5 µl/Test,right).