PE Rabbit anti-Human CD153 mAb

www.abclonal.com

ABclonal

Catalog No.: A25121

Basic Information

Observed MW

Calculated MW

26kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC64953-PE

Conjugate

PE. Ex:565nm. Em:574nm.

Background

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF8/CD30, which is a cell surface antigen and a marker for Hodgkin lymphoma and related hematologic malignancies. The engagement of this cytokine expressed on B cell surface plays an inhibitory role in modulating Ig class switch. This cytokine was shown to enhance cell proliferation of some lymphoma cell lines, while to induce cell death and reduce cell proliferation of other lymphoma cell lines. The pleiotropic biologic activities of this cytokine on different CD30+lymphoma cell lines may play a pathophysiologic role in Hodgkin's and some non-Hodgkin's lymphomas. Two transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Immunogen Information

Gene ID 944 **Swiss Prot**

P32971

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 63-234 of human CD153 (NP_001235.1).

Synonyms

CD153; CD30L; CD30LG; TNLG3A

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

Product Information


SourceIsotypePurificationRabbitIgGAffinity purification


Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data





Flow cytometry: 1X10^6 K-562 cells (negative control,left) and Daudi cells (right) were surface-stained with PE Rabbit anti-Human CD153 mAb (A25121,5 $\,\mu$ l/Test,orange line) or PE Rabbit IgG isotype control (5 $\,\mu$ l/Test,blue line). Nonfluorescently stained cells were used as blank control (red line).

Flow cytometry: $1X10^6$ Daudi cells were surface-stained with PE Rabbit IgG isotype control (5 μ I/Test,left) or PE Rabbit anti-Human CD153 mAb (A25121,5 μ I/Test,right).