

KPNA6 Rabbit PolymAb®

Catalog No.: A25199

Basic Information

Observed MW

60kDa/

Calculated MW

60kDa

Category

Primary antibody

Applications

ELISA,WB,IHC-P,IF/ICC

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC64379_ARC64377

Background

Nucleocytoplasmic transport, a signal- and energy-dependent process, takes place through nuclear pore complexes embedded in the nuclear envelope. The import of proteins containing a nuclear localization signal (NLS) requires the NLS import receptor, a heterodimer of importin alpha and beta subunits also known as karyopherins. Importin alpha binds the NLS-containing cargo in the cytoplasm and importin beta docks the complex at the cytoplasmic side of the nuclear pore complex. In the presence of nucleoside triphosphates and the small GTP binding protein Ran, the complex moves into the nuclear pore complex and the importin subunits dissociate. Importin alpha enters the nucleoplasm with its passenger protein and importin beta remains at the pore. The protein encoded by this gene is a member of the importin alpha family.

Recommended Dilutions

WB	1:3000 - 1:20000
IHC-P	1:1000-1:5000
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

23633

Swiss Prot

O60684

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 69-169 of human KPNA6 (NP_036448.1).

Synonyms

IPOA7; KPNA6

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from HeLa cells using KPNA6 Rabbit PolymAb® (A25199) at 1:20000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 20s.




Confocal imaging of A-431 cells using KPNA6 Rabbit PolymAb® (A25199, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of HeLa cells using KPNA6 Rabbit PolymAb® (A25199, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Confocal imaging of NIH/3T3 cells using KPNA6 Rabbit PolymAb® (A25199, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunohistochemistry analysis of KPNA6 in paraffin-embedded human breast cancer tissue using KPNA6 Rabbit PolymAb® (A25199) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of KPNA6 in paraffin-embedded human esophagus tissue using KPNA6 Rabbit PolymAb® (A25199) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of KPNA6 in paraffin-embedded rat colon tissue using KPNA6 Rabbit PolymAb® (A25199) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.


Validation Data




Immunohistochemistry analysis of KPNA6 in paraffin-embedded mouse colon tissue using KPNA6 Rabbit PolymAb® (A25199) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of KPNA6 in paraffin-embedded human breast cancer tissue using KPNA6 Rabbit PolymAb® (A25199) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of KPNA6 in paraffin-embedded rat colon tissue using KPNA6 Rabbit PolymAb® (A25199) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of KPNA6 in paraffin-embedded human esophagus tissue using KPNA6 Rabbit PolymAb® (A25199) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of KPNA6 in paraffin-embedded mouse colon tissue using KPNA6 Rabbit PolymAb® (A25199) at a dilution of 1:3000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.