

CREBBP Rabbit mAb

Catalog No.: A25323 **Recombinant**

Basic Information

Observed MW

300kDa

Calculated MW

265kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P, IF/ICC, ChIP

Cross-Reactivity

Human, Mouse, Rat

CloneNo number

ARC3258

Background

This gene is ubiquitously expressed and is involved in the transcriptional coactivation of many different transcription factors. First isolated as a nuclear protein that binds to cAMP-response element binding protein (CREB), this gene is now known to play critical roles in embryonic development, growth control, and homeostasis by coupling chromatin remodeling to transcription factor recognition. The protein encoded by this gene has intrinsic histone acetyltransferase activity and also acts as a scaffold to stabilize additional protein interactions with the transcription complex. This protein acetylates both histone and non-histone proteins. This protein shares regions of very high sequence similarity with protein p300 in its bromodomain, cysteine-histidine-rich regions, and histone acetyltransferase domain. Mutations in this gene cause Rubinstein-Taybi syndrome (RTS). Chromosomal translocations involving this gene have been associated with acute myeloid leukemia. Alternative splicing results in multiple transcript variants encoding different isoforms.

Recommended Dilutions

WB 1:500 - 1:1000**IHC-P** 1:500 - 1:2000**IF/ICC** 1:50 - 1:200**ChIP** 3µg antibody for
10µg-15µg of Chromatin

Immunogen Information

Gene ID

1387

Swiss Prot

Q92793

Immunogen

Synthetic peptide

Synonyms

CBP; RSTS; KAT3A; MKHK1; RSTS1; CREBBP

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of lysates from HeLa cells using CREBBP Rabbit mAb(A25323) at 1:1000 dilution. Secondary antibody:HRP Goat Anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection:ECL Basic Kit (RM00020). Exposure time:45s.




Chromatin immunoprecipitation was performed with 15 µg of cross-linked chromatin from HeLa cells, using 5 µg of CREBBP (A25323) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.




Chromatin immunoprecipitation was performed with 15 µg of cross-linked chromatin from HepG2 cells, using 5 µg of CREBBP (A25323) and Rabbit IgG isotype control (AC042). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.



Confocal imaging of HeLa cells using CREBBP Rabbit mAb (A25323, dilution 1:500) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500).




Confocal imaging of NIH/3T3 cells using CREBBP Rabbit mAb (A25323, dilution 1:500) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500).



Immunohistochemistry analysis of CREBBP in paraffin-embedded human spleen tissue using CREBBP Rabbit mAb (A25323) at a dilution of 1:2000 (40x lens). High pressure liquid chromatography (HPLC) analysis of the antibody showed a purity of >95%.


Validation Data

1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunohistochemistry analysis of CREBBP in paraffin-embedded human breast cancer tissue using CREBBP Rabbit mAb (A25323) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

dilution 1:500) (Red). The cells were counterstained with α -Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunohistochemistry analysis of CREBBP in paraffin-embedded mouse brain tissue using CREBBP Rabbit mAb (A25323) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of CREBBP in paraffin-embedded rat liver tissue using CREBBP Rabbit mAb (A25323) at a dilution of 1:2000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.