ABclonal

www.abclonal.com

ABflo® 488 Rabbit anti-Human CD24 mAb

Catalog No.: A25380

Basic Information

Observed MW

Refer to figures

Calculated MW

8kDa; 13kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC66763-ABflo488

Conjugate

ABflo® 488. Ex:491nm. Em:516nm.

Background

This gene encodes a sialoglycoprotein that is expressed on mature granulocytes and B cells and modulates growth and differentiation signals to these cells. The precursor protein is cleaved to a short 32 amino acid mature peptide which is anchored via a glycosyl phosphatidylinositol (GPI) link to the cell surface. This gene was missing from previous genome assemblies, but is properly located on chromosome 6. Non-transcribed pseudogenes have been designated on chromosomes 1, 15, 20, and Y. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

FC

5 μl per 10^6 cells in 100 μl volume

Immunogen Information

Gene ID 100133941

Swiss Prot

41 P25063

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-80 of human CD24 (NP_037362.1).

Synonyms

CD24A; CD24

Contact

a		400-999-6126
\bowtie		cn.market@abclonal.com.cn
\odot	T	www.abclonal.com.cn

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage


Store at 2-8°C. Avoid freeze.


Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data









Flow cytometry: 1X10^6 293F cells (Low Expression,left) and MCF7 cells (right) were surface-stained with ABflo® 488 Rabbit anti-Human CD24 mAb (A25380,5 μ I/Test,orange line) or ABflo® 488 Rabbit IgG isotype control (A22069,5 μ I/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10^6 MCF7 cells were surface-stained with ABflo® 488 Rabbit IgG isotype control (A22069,5 µl/Test,left) or ABflo® 488 Rabbit anti-Human CD24 mAb (A25380,5 µl/Test,right).

Confocal imaging of MCF7 cells using ABflo® 488 Rabbit anti-Human CD24 mAb (A25380, dilution 1:100). DAPI was used for nuclear staining (Blue). Objective: 100x.