Leader in Biomolecular Solutions for Life Science

# **ARHGEF2** Rabbit pAb

Catalog No.: A2539



### **Basic Information**

Observed MW 120kDa

Calculated MW 112kDa

Category Primary antibody

Applications WB,ELISA

Cross-Reactivity Human

### Background

Rho GTPases play a fundamental role in numerous cellular processes that are initiated by extracellular stimuli that work through G protein coupled receptors. The encoded protein may form complex with G proteins and stimulate rho-dependent signals. Alternatively spliced transcript variants encoding different isoforms have been identified.

### **Recommended Dilutions**

| WB    | 1:500 - 1:1000   |
|-------|--|
| ELISA | Recommended starting<br>concentration is 1 µg/mL.<br>Please optimize the<br>concentration based on<br>your specific assay<br>requirements. |

## **Immunogen Information**

#### **Gene ID** 9181

Swiss Prot Q92974

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 850-950 of human ARHGEF2 (NP\_001155855.1).

Synonyms

GEF; Lfc; P40; GEFH1; LFP40; GEF-H1; NEDMHM; ARHGEF2

### Contact


| 6        | 400-999-6126              |
|----------|---------------------------|
| $\times$ | cn.market@abclonal.com.cn |
| €        | www.abclonal.com.cn       |

# **Product Information**

**Source** Rabbit **Isotype** IgG Purification Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using ARHGEF2 Rabbit pAb (A2539) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 180s.