Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Estrogen Receptor beta Rabbit pAb

Catalog No.: A2546SP **11 Publications**

Basic Information

Observed MW
50-60 kDa

Calculated MW
35-59 kDa

Category
Primary antibody

Applications
WB,IP,IF/ICC,IF-P,ELISA

Cross-Reactivity
Human, Mouse, Rat

Background

This gene encodes a member of the family of estrogen receptors and superfamily of nuclear receptor transcription factors. The gene product contains an N-terminal DNA binding domain and C-terminal ligand binding domain and is localized to the nucleus, cytoplasm, and mitochondria. Upon binding to 17beta-estradiol or related ligands, the encoded protein forms homo- or hetero-dimers that interact with specific DNA sequences to activate transcription. Some isoforms dominantly inhibit the activity of other estrogen receptor family members. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been fully characterized.

Recommended Dilutions

WB	1:1000 - 1:2000
IP	0.5 μg - 4 μg antibody for 200 μg - 400 μg extracts of whole cells
IF/ICC	1:200 - 1:400
IF-P	1:200 - 1:400
ELISA	Recommended starting concentration is 1 μg/mL. Please optimize the concentration based on your specific assay requirements.For high-ratio antibody dilutions (≥1:10000)，a sequential dilution method is strongly recommended to ensure measurement accuracy.

Immunogen Information

Gene ID
2100

Swiss Prot
Q92731

Immunogen
Recombinant protein (or fragment).This information is considered to be commercially sensitive.

Synonyms
Erb; ESRB; ODG8; ESTRB; NR3A2; ER-BETA; ESR-BETA

Product Information

Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20℃


Contact

400-999-6126


cn.market@abclonal.com.cn

www.abclonal.com.cn


Validation Data




Western blot analysis of lysates from Rat uterus using Estrogen Receptor beta Rabbit pAb (A2546SP) at 1:2000 dilution incubated overnight at 4℃.|Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.|Lysates/proteins: 25 µg per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit (RM00020).|Exposure time: 30 s



Confocal imaging of MCF7 cells using Estrogen Receptor beta Rabbit pAb (A2546SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x



Confocal imaging of NIH/3T3 cells using Estrogen Receptor beta Rabbit pAb (A2546SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x



Confocal imaging of paraffin-embedded Rat uterus using Estrogen Receptor beta Rabbit pAb (A2546SP, dilution 1:200) followed by a further incubation with Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). DAPI was used for nuclear staining (Blue). High pressure antigen retrieval performed with 0.01M Citrate Buffer (pH 6.0) prior to IF staining. Objective: 40x