

FITC Rabbit Anti-Goat IgG (Fc) fragment mAb

Catalog No.: A25513

Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

FC

Cross-Reactivity

Goat

CloneNo number

ARC56943-FITC

Conjugate

FITC. Ex:491nm. Em:516nm.

Background

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Immunogen Information

Gene ID


Swiss Prot

Immunogen

Recombinant protein of goat IgG (Fc) fragment.

Synonyms

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification



Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data



Flow cytometry: 1×10^6 Jurkat cells (negative control, left) and A-431 cells (right) were surface-stained with Goat anti-Human TROP-2 mAb (IgG, κ , orange line) or FITC Rabbit IgG isotype control (5 μ l/Test, blue line), followed by FITC Rabbit Anti-Goat IgG (Fc) fragment mAb (A25513, 5 μ l/Test) staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 A-431 cells were surface-stained with FITC Rabbit IgG isotype control (5 μ l/Test, left) or Goat anti-Human TROP-2 mAb (IgG, κ , right). FITC Rabbit Anti-Goat IgG (Fc) fragment mAb (A25513, 5 μ l/Test) was used as a secondary antibody.