

ABflo® 700 Rabbit Anti-Goat IgG (Fc) fragment mAb

Catalog No.: A25515

Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

FC

Cross-Reactivity

Goat

CloneNo number

ARC56943-ABflo700

Conjugate

ABflo® 700. Ex:690nm. Em:713nm.

Background

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Recombinant protein of goat IgG (Fc) fragment.

Synonyms

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data



Flow cytometry: 1×10^6 Jurkat cells (negative control, left) and A-431 cells (right) were surface-stained with Goat anti-Human TROP-2 mAb (IgG, κ, orange line) or ABflo® 700 Rabbit IgG isotype control (5 μl/Test, blue line), followed by ABflo® 700 Rabbit Anti-Goat IgG (Fc) fragment mAb (A25515, 5 μl/Test) staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 A-431 cells were surface-stained with ABflo® 700 Rabbit IgG isotype control (5 μl/Test, left) or Goat anti-Human TROP-2 mAb (IgG, κ, right). ABflo® 700 Rabbit Anti-Goat IgG (Fc) fragment mAb (A25515, 5 μl/Test) was used as a secondary antibody.