

ABflo® 700 Rabbit Anti-Goat IgG (Fc) fragment mAb

Catalog No.: A25515

Basic Information

Observed MW

Calculated MW

Category

Primary antibody

Applications

FC

Cross-Reactivity

Goat

CloneNo number

ARC56943

Conjugate

ABflo® 700. Ex:690nm. Em:713nm.

Background

Recommended Dilutions

FC 5 µl per 10⁶ cells in
 100 µl volume

Immunogen Information

Gene ID

Swiss Prot

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification



Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Flow cytometry: 1X10⁶ Jurkat cells (negative control, left) and A-431 cells (right) were surface-stained with Goat anti-Human TROP-2 mAb (IgG, κ, orange line) or ABflo® 700 Rabbit IgG isotype control (5 μl/Test, blue line), followed by ABflo® 700 Rabbit Anti-Goat IgG (Fc) fragment mAb (A25515, 5 μl/Test) staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1X10⁶ A-431 cells were surface-stained with ABflo® 700 Rabbit IgG isotype control (5 μl/Test, left) or Goat anti-Human TROP-2 mAb (IgG, κ, right). ABflo® 700 Rabbit Anti-Goat IgG (Fc) fragment mAb (A25515, 5 μl/Test) was used as a secondary antibody.