

# PE Rabbit Anti-Goat IgG (Fc) fragment mAb

Catalog No.: A25517

## Basic Information

**Observed MW**

**Calculated MW**

**Category**

Primary antibody

**Applications**

FC

**Cross-Reactivity**

Goat

**CloneNo number**

ARC56943-PE

**Conjugate**

PE. Ex:565nm. Em:574nm.

## Background

## Recommended Dilutions

**FC**                    5  $\mu$ l per  $10^6$  cells in  
                                 100  $\mu$ l volume

## Immunogen Information

**Gene ID**

**Swiss Prot**

**Immunogen**

Recombinant protein of goat IgG (Fc) fragment.

**Synonyms**

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

## Validation Data



Flow cytometry:  $1 \times 10^6$  Jurkat cells (negative control, left) and A-431 cells (right) were surface-stained with Goat anti-Human TROP-2 mAb (IgG,  $\kappa$ , orange line) or PE Rabbit IgG isotype control (A24172, 5  $\mu$ l/Test, blue line), followed by PE Rabbit Anti-Goat IgG (Fc) fragment mAb (A25517, 5  $\mu$ l/Test) staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1 \times 10^6$  A-431 cells were surface-stained with PE Rabbit IgG isotype control (A24172, 5  $\mu$ l/Test, left) or Goat anti-Human TROP-2 mAb (IgG,  $\kappa$ , right). PE Rabbit Anti-Goat IgG (Fc) fragment mAb (A25517, 5  $\mu$ l/Test) was used as a secondary antibody.