

# Galectin 3/LGALS3 Rabbit mAb

Catalog No.: A25538 **Recombinant**

## Basic Information

### Observed MW

28kDa

### Calculated MW

26kDa

### Category

Primary antibody

### Applications

WB,IF/ICC,IP,FC,ELISA

### Cross-Reactivity

Human, Mouse

### CloneNo number

ARC58285

## Recommended Dilutions

**WB** 1:9000 - 1:54000

**IF/ICC** 1:200-1:800

**IP** 0.5µg-4µg antibody for  
200µg-400µg extracts of  
whole cells

**FC** 1:100 - 1:500

**ELISA** Recommended starting  
concentration is 1 µg/mL.  
Please optimize the  
concentration based on  
your specific assay  
requirements.

## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

This gene encodes a member of the galectin family of carbohydrate binding proteins. Members of this protein family have an affinity for beta-galactosides. The encoded protein is characterized by an N-terminal proline-rich tandem repeat domain and a single C-terminal carbohydrate recognition domain. This protein can self-associate through the N-terminal domain allowing it to bind to multivalent saccharide ligands. This protein localizes to the extracellular matrix, the cytoplasm and the nucleus. This protein plays a role in numerous cellular functions including apoptosis, innate immunity, cell adhesion and T-cell regulation. The protein exhibits antimicrobial activity against bacteria and fungi. Alternate splicing results in multiple transcript variants.

## Immunogen Information

### Gene ID

3958

### Swiss Prot

P17931

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human Galectin 3/LGALS3 (NP\_002297.2).

### Synonyms

L31; GAL3; MAC2; CBP35; GALBP; GALIG; LGALS2

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.


## Validation Data




Immunoprecipitation of Galectin 3/LGALS3 from 300 µg extracts of HeLa cells was performed using 3 µg of Galectin 3/LGALS3 Rabbit mAb (A25538). Rabbit IgG isotype control (AC042) was used to precipitate the Control IgG sample. IP samples were eluted with 1X reducing Laemmli Buffer. The Input lane represents 10% of the total input. Western blot analysis of immunoprecipitates was conducted using Galectin 3/LGALS3 Rabbit mAb (A25538) at 1:10000 dilution.




Western blot analysis of various lysates using Galectin 3/LGALS3 Rabbit mAb (A25538) at 1:9000 dilution incubated overnight at 4°C.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25 µg per lane.  
 Blocking buffer: 3 % nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 10s.



Flow cytometry:  $1 \times 10^6$  Jurkat cells (negative control, left) and MCF7 cells (right) were surface-stained with Galectin 3/LGALS3 Rabbit mAb (A25538, 2 µg/mL, orange line) or Rabbit IgG isotype control (AC042, 2 µg/mL, blue line), followed by FITC conjugated goat anti-Rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry:  $1 \times 10^6$  MCF7 cells were surface-stained with Rabbit IgG isotype control (AC042, 2 µg/mL, left) or Galectin 3/LGALS3 Rabbit mAb (A25538, 2 µg/mL, right), followed by FITC conjugated goat anti-Rabbit pAb staining.



Immunofluorescence analysis of HeLa cells using Galectin 3/LGALS3 Rabbit mAb (A25538) at a dilution of 1:400 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.