

PerCP Rabbit anti-Human/Monkey CD45 mAb

Catalog No.: A25624

Basic Information

Observed MW**Calculated MW**

131kDa/136kDa/141kDa/143kDa/147kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Human, Cynomolgus

CloneNo number

ARC5024

Conjugate

PerCP. Ex:482nm. Em:678nm.

Background

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Immunogen Information

Gene ID

5788

Swiss Prot

P08575

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

LCA; LY5; B220; CD45; L-CA; T200; CD45R; GP180; IMD105

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Validation Data




Flow cytometry: 1×10^6 293T cells (negative control, left) and Jurkat cells (right) were surface-stained with PerCP Rabbit anti-Human CD45 mAb (A25624, 5 μ l/Test, orange line) or PerCP Rabbit IgG isotype control (A24204, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).



Flow cytometry: 1×10^6 Jurkat cells were surface-stained with PerCP Rabbit IgG isotype control (A24204, 5 μ l/Test, left) or PerCP Rabbit anti-Human CD45 mAb (A25624, 5 μ l/Test, right).



Flow cytometry: 1×10^6 Human PBMC were surface-stained with PerCP Rabbit anti-Human CD45 mAb (A25624, 5 μ l/Test, orange line) or PerCP Rabbit IgG isotype control (5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line). Cells in the lymphocyte gate were used for analysis.



Flow cytometry: 1×10^6 Human PBMC were surface-stained with PerCP Rabbit IgG isotype control (5 μ l/Test, left) or PerCP Rabbit anti-Human CD45 mAb (A25624, 5 μ l/Test, right). Cells in the lymphocyte gate were used for analysis.