

# PerCP Rabbit anti-Human CD45 mAb

Catalog No.: A25624

## Basic Information

### Observed MW

### Calculated MW

131kDa/136kDa/141kDa/143kDa/147kDa

### Category

Primary antibody

### Applications

FC

### Cross-Reactivity

Human

### CloneNo number

ARC5024-PerCP

### Conjugate

PerCP. Ex:482nm. Em:678nm.

## Recommended Dilutions

FC 5  $\mu$ l per  $10^6$  cells in  
100  $\mu$ l volume

## Background

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported.

## Immunogen Information

### Gene ID

5788

### Swiss Prot

P08575

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 26-577 of human CD45 (P08575).

### Synonyms

LCA; LY5; B220; CD45; L-CA; T200; CD45R; GP180; IMD105

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

## Validation Data



Flow cytometry:  $1 \times 10^6$  293T cells (negative control, left) and Jurkat cells (right) were surface-stained with PerCP Rabbit anti-Human CD45 mAb (A25624, 5  $\mu$ l/Test, orange line) or PerCP Rabbit IgG isotype control (A24204, 5  $\mu$ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1 \times 10^6$  Jurkat cells were surface-stained with PerCP Rabbit IgG isotype control (A24204, 5  $\mu$ l/Test, left) or PerCP Rabbit anti-Human CD45 mAb (A25624, 5  $\mu$ l/Test, right).