

# [KO Validated] LAMP2 Rabbit mAb

Catalog No.: A25646 **KO** **Validated** **Recombinant**

## Basic Information

### Observed MW

100-130kDa

### Calculated MW

45kDa

### Category

Primary antibody

### Applications

ELISA,WB,IHC-P,IF/ICC,FC (intra)

### Cross-Reactivity

Human

### CloneNo number

ARC54767

## Background

The protein encoded by this gene is a member of a family of membrane glycoproteins. This glycoprotein provides selectins with carbohydrate ligands. It may play a role in tumor cell metastasis. It may also function in the protection, maintenance, and adhesion of the lysosome. Alternative splicing of this gene results in multiple transcript variants encoding distinct proteins.

## Recommended Dilutions

<b>WB</b>	1:10000 - 1:20000
<b>IHC</b>	1:50 - 1:200
<b>IF/ICC</b>	1:50 - 1:200
<b>FC (intra)</b>	1:500 - 1:1000

## Immunogen Information

### Gene ID

3920

### Swiss Prot

P13473

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 29-375 of human LAMP2 (NP\_002285.1).

### Synonyms

DND; LAMPB; CD107b; LAMP-2; LGP-96; LGP110

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.


## Validation Data




Western blot analysis of lysates from A549 cells using [KO Validated] LAMP2 Rabbit mAb (A25646) at 1:20000 dilution.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25 µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time:10s.




Western blot analysis of lysates from wild type (WT) and LAMP2 knockout (KO) HeLa cells using [KO Validated] LAMP2 Rabbit mAb (A25646) at 1:20000 dilution.  
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
 Lysates/proteins: 25 µg per lane.  
 Blocking buffer: 3% nonfat dry milk in TBST.  
 Detection: ECL Basic Kit (RM00020).  
 Exposure time: 10s.



Flow cytometry:  $1 \times 10^6$  knockout (KO) HeLa cells (negative control, left) and HeLa cells (right) were intracellularly-stained with [KO Validated] LAMP2 Rabbit mAb (A25646, 2 µg/mL, orange line) or Rabbit IgG isotype control (AC042, 2 µg/mL, blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).




Flow cytometry:  $1 \times 10^6$  knockout (KO) HeLa cells (negative control, left) and HeLa cells (right) were intracellularly-stained with Rabbit IgG isotype control (AC042, 2 µg/mL, left) or [KO Validated] LAMP2 Rabbit mAb (A25646, 2 µg/mL, right), followed by Alexa Fluor® 647 conjugated goat anti-Rabbit pAb staining.



Confocal imaging of HeLa cells using [KO Validated] LAMP2 Rabbit mAb (A25646, dilution 1:200) followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) (AS007, dilution 1:500) (Red). The cells were counterstained with α-Tubulin Mouse mAb (AC012, dilution 1:400) followed by incubation with ABflo® 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (AS076, dilution 1:500) (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.

## Validation Data

---



Immunohistochemistry analysis of paraffin-embedded Human kidney tissue using [KO Validated] LAMP2 Rabbit mAb (A25646) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.