# PE Rabbit anti-Human CD124/IL-4Rα mAb

ABclonal www.abclonal.com

Catalog No.: A25725

### **Basic Information**

#### **Observed MW**

# Calculated MW

26kDa/88kDa/90kDa

## Category

Primary antibody

### **Applications**

FC

### **Cross-Reactivity**

Human

#### CloneNo number

ARC66266-PE

### Conjugate

PE. Ex:565nm. Em:574nm.

# **Background**

This gene encodes the alpha chain of the interleukin-4 receptor, a type I transmembrane protein that can bind interleukin 4 and interleukin 13 to regulate IgE production. The encoded protein also can bind interleukin 4 to promote differentiation of Th2 cells. A soluble form of the encoded protein can be produced by proteolysis of the membrane-bound protein, and this soluble form can inhibit IL4-mediated cell proliferation and IL5 upregulation by T-cells. Allelic variations in this gene have been associated with atopy, a condition that can manifest itself as allergic rhinitis, sinusitus, asthma, or eczema. Polymorphisms in this gene are also associated with resistance to human immunodeficiency virus type-1 infection. Alternate splicing results in multiple transcript variants.

## **Recommended Dilutions**

FC

5 μl per 10^6 cells in 100 μl volume

# Immunogen Information

Gene ID 3566 **Swiss Prot** 

P24394

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 24-232 of human CD124/IL-4R $\alpha$  (NP\_000409.1).

### **Synonyms**

CD124; IL4RA; IL-4RA

## **Contact**

2		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	T	www.abclonal.com.cn

### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### **Storage**


Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

# **Validation Data**







Flow cytometry: 1X10^6 MOLT-4 cells (negative control,left) and Daudi cells (right) were surface-stained with PE Rabbit anti-Human CD124/IL-4R $\alpha$  mAb (A25725,5  $\mu$ I/Test,orange line) or PE Rabbit IgG isotype control (A24172,5  $\mu$ I/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1X10^6$  Daudi cells were surface-stained with PE Rabbit IgG isotype control (A24172,5  $\mu$ I/Test,Ieft) or PE Rabbit anti-Human CD124/IL-4R $\alpha$  mAb (A25725,5  $\mu$ I/Test,right).