

APC Rabbit anti-Mouse CD366 (TIM-3) mAb

Catalog No.: A25726

Basic Information

Observed MW

Refer to figures

Calculated MW

22kDa/31kDa

Category

Primary antibody

Applications

FC

Cross-Reactivity

Mouse

CloneNo number

ARC59425

Conjugate

APC. Ex:650nm. Em:660nm.

Recommended Dilutions

FC 5 μ l per 10^6 cells in
100 μ l volume

Background

Predicted to enable metal ion binding activity. Involved in several processes, including regulation of cytokine production; regulation of leukocyte activation; and toll-like receptor signaling pathway. Located in cell surface; early endosome; and immunological synapse. Is expressed in several structures, including dorsal aorta; gut; liver; lung; and reproductive system. Human ortholog(s) of this gene implicated in hepatocellular carcinoma and renal cell carcinoma. Orthologous to human HAVCR2 (hepatitis A virus cellular receptor 2).

Immunogen Information

Gene ID

171285

Swiss Prot

Q8VIM0

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

Tim3; TIM-3; Timd3

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Flow cytometry: 1×10^6 NIH/3T3 cells (Low Expression, left) and RAW 264.7 cells (right) were surface-stained with APC Rabbit anti-Mouse CD366 (TIM-3) mAb (A25726, 5 μ l/Test, orange line) or APC Rabbit IgG isotype control (A24173, 5 μ l/Test, blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 RAW 264.7 cells were surface-stained with APC Rabbit IgG isotype control (A24173, 5 μ l/Test, left) or APC Rabbit anti-Mouse CD366 (TIM-3) mAb (A25726, 5 μ l/Test, right).