# PE Rabbit anti-Mouse IL1R1/CD121a mAb

ABclonal www.abclonal.com

Catalog No.: A25858

#### **Basic Information**

#### **Observed MW**

**Calculated MW** 

67kDa

Category

Primary antibody

**Applications** 

FC

**Cross-Reactivity** 

Mouse

CloneNo number

ARC65805-PE

Conjugate

PE. Ex:565nm. Em:574nm.

## **Background**

Enables interleukin-1 binding activity; interleukin-1 receptor activity; and protease binding activity. Involved in interleukin-1-mediated signaling pathway. Acts upstream of or within several processes, including cytokine-mediated signaling pathway; positive regulation of interleukin-1-mediated signaling pathway; and positive regulation of neutrophil extravasation. Located in external side of plasma membrane. Is expressed in brain; dorsal aorta; liver; reproductive system; and spleen. Used to study type 1 diabetes mellitus. Human ortholog(s) of this gene implicated in systemic scleroderma and type 1 diabetes mellitus. Orthologous to human IL1R1 (interleukin 1 receptor type 1).

### **Recommended Dilutions**

FC

5  $\mu$ l per 10^6 cells in 100  $\mu$ l volume

## Immunogen Information

**Gene ID** 16177

**Swiss Prot** 

P13504

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 20-338 of mouse IL1R1/CD121a (NP\_032388.1).

#### **Synonyms**

IL-iR; CD121a; CD121b; IL-1R1; II1r-1; IL-1R-1; IL-1RT1; IL-1RT-1; IL-1R-alpha

#### **Contact**

<b>a</b>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
$\odot$	Т	www.abclonal.com.cn

#### **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

#### **Storage**

Store at 2-8°C. Avoid freeze.

Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

## **Validation Data**









Flow cytometry: 1X10^6 Neuro-2a cells (negative control,left) and NIH/3T3 cells (right) were surface-stained with PE Rabbit anti-Mouse IL1R1/CD121a mAb (A25858,5 µl/Test,orange line) or PE Rabbit IgG isotype control (A24172,5 µl/Test,blue line). Nonfluorescently stained cells were used as blank control (red line).

Flow cytometry:  $1X10^6$  NIH/3T3 cells were surface-stained with PE Rabbit IgG isotype control (A24172,5  $\mu$ I/Test,left) or PE Rabbit anti-Mouse IL1R1/CD121a mAb (A25858,5  $\mu$ I/Test,right).