

Nanog Rabbit mAb

Catalog No.: A25887 **Recombinant**

Basic Information

Observed MW

35-42kDa

Calculated MW

34kDa

Category

Primary antibody

Applications

ELISA, WB, ChIP, FC (intra)

Cross-Reactivity

Human, Mouse

CloneNo number

ARC62770

Background

The protein encoded by this gene is a DNA binding homeobox transcription factor involved in embryonic stem (ES) cell proliferation, renewal, and pluripotency. The encoded protein can block ES cell differentiation and can also autorepress its own expression in differentiating cells. Several transcript variants encoding different isoforms have been found for this gene.

Recommended Dilutions

WB 1:500 - 1:1000

ChIP 5µg antibody for
10µg-15µg of Chromatin

FC (intra) 1:500 - 1:1000

Immunogen Information

Gene ID

71950

Swiss Prot

Q80Z64

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

ENK; Stm1; ecat4; 2410002E02Rik

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.




Validation Data



Western blot analysis of lysates from NCCIT cells using Nanog Rabbit mAb (A25887) at 1:1000 dilution incubated overnight at 4°C.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25 µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 90s.



Chromatin immunoprecipitation was performed with 10 µg of cross-linked chromatin from F9 cells, using 5 µg of Nanog Rabbit mAb (A25887) and Rabbit Control IgG (AC005). The enrichment of immunoprecipitated DNA at different genomic loci was examined by quantitative PCR. The histogram compares the ratio of the immunoprecipitated DNA to the input at given loci.




Flow cytometry: 1×10^6 HeLa cells (negative control, left) and NTERA-2 cells (right) were intracellularly-stained with Nanog Rabbit mAb (A25887, 2 µg/mL, orange line) or Rabbit IgG isotype control (AC042, 2 µg/mL, blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry: 1×10^6 NTERA-2 cells were intracellularly-stained with Rabbit IgG isotype control (AC042, 2 µg/mL, left) or Nanog Rabbit mAb (A25887, 2 µg/mL, right), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining.

Flow cytometry: 1×10^6 NIH/3T3 cells (negative control, left) and F9 cells (right) were intracellularly-stained with Nanog Rabbit mAb (A25887, 2 µg/mL, orange line) or Rabbit IgG isotype control (AC042, 2 µg/mL, blue line), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining. Non-fluorescently stained cells were used as blank control (red line).

Validation Data



Flow cytometry: 1×10^6 F9 cells were intracellularly-stained with Rabbit IgG isotype control (AC042,2 $\mu\text{g}/\text{mL}$, left) or Nanog Rabbit mAb (A25887,2 $\mu\text{g}/\text{mL}$, right), followed by Alexa Fluor® 647 conjugated goat anti-rabbit pAb staining.