

ABflo® 594 Rabbit anti-Human/Monkey CD69 mAb

Catalog No.: A25893

Basic Information

Observed MW

Calculated MW
23kDa

Category
Primary antibody

Applications
FC

Cross-Reactivity
Human, Cynomolgus

CloneNo number
ARC68505

Conjugate
ABflo® 594. Ex:588nm. Em:604nm.

Background

This gene encodes a member of the calcium dependent lectin superfamily of type II transmembrane receptors. Expression of the encoded protein is induced upon activation of T lymphocytes, and may play a role in proliferation. Furthermore, the protein may act to transmit signals in natural killer cells and platelets.

Recommended Dilutions

FC 5 µl per 10⁶ cells in
100 µl volume

Immunogen Information

Gene ID	Swiss Prot
969	Q07108

Immunogen

Recombinant protein (or fragment). This information is considered to be commercially sensitive.

Synonyms

AIM; EA1; MLR-3; CLEC2C; GP32/28; BL-AC/P26

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information


Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage


Store at 2-8°C. Avoid freeze.

Buffer: PBS containing 0.2% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Validation Data



Flow cytometry: 1×10^6 Cynomolgus PBMC (untreated, left) and Cynomolgus PBMC (treated with 50 ng/mL PMA and 1 µg/mL Ionomycin for 6 hours, right) were surface-stained with ABflo® 450 Rabbit anti-Human/Monkey CD3 mAb (A27177, 5 µl/Test) and ABflo® 594 Rabbit anti-Human/Monkey CD69 mAb (A25893, 5 µl/Test). Cells in the lymphocyte gate were used for analysis.



Flow cytometry: 1×10^6 Human PBMC (untreated, left) and Human PBMC (treated with 50 ng/mL PMA and 1 µg/mL Ionomycin for 6 hours, right) were surface-stained with ABflo® 594 Rabbit anti-Human/Monkey CD69 mAb (A25893, 5 µl/Test). Cells in the lymphocyte gate were used for analysis.